


## Volatile Organic Compound Emissions from Humans Indoors

Xiaochen Tang,<sup>†</sup> Pawel K. Misztal,<sup>‡</sup> William W Nazaroff,<sup>†</sup> and Allen H. Goldstein<sup>\*,†,‡</sup><sup>†</sup>Department of Civil and Environmental Engineering, University of California, Berkeley, California 94720-1710 United States<sup>‡</sup>Department of Environmental Science, Policy and Management, University of California, Berkeley, California 94720-3114 United States

## Supporting Information

**ABSTRACT:** Research on the sources of indoor airborne chemicals has traditionally focused on outdoor air, building materials, furnishings, and activities such as smoking, cooking, and cleaning. Relatively little research has examined the direct role of occupant emissions, even though this source clearly contributes to indoor volatile organic compounds (VOCs) and influences indoor chemistry. In this work, we quantify occupant-related gaseous VOC emissions in a university classroom using a proton-transfer-reaction time-of-flight mass spectrometer. Time-resolved concentrations of VOCs in room air and supply air were measured continuously during occupied and unoccupied periods. The emission factor for each human-emitted VOC was determined by dividing the occupant-associated source rate by the corresponding occupancy. Among the most abundant species detected were compounds associated with personal care products. Also prominent were human metabolic emissions, such as isoprene, methanol, acetone, and acetic acid. Additional sources included human skin oil oxidation by ozone, producing compounds such as 4-oxopentanal (4-OPA) and 6-methyl-5-hepten-2-one (6-MHO). By mass, human-emitted VOCs were the dominant source (57%) during occupied periods in a well-ventilated classroom, with ventilation supply air the second most important (35%), and indoor nonoccupant emissions the least (8%). The total occupant-associated VOC emission factor was 6.3 mg h<sup>-1</sup> per person.



## INTRODUCTION

Human emissions of volatile organic compounds (VOCs) can strongly influence indoor air quality. Since humans spend most of their time indoors, most air inhaled by people is indoor air, and, therefore, occupant emissions of VOCs must affect humankind's aggregate inhalation exposure. Yet, notwithstanding its significance, remarkably little research has focused on characterizing occupant-associated VOC emissions to indoor environments.

Historically, ventilation rates in buildings were set at levels designed to control the perceived odors associated with human occupants.<sup>1</sup> Although there have been changes over time, human perception and subjective assessment of the acceptability of indoor air remains an important basis for current ventilation standards and practice.<sup>2</sup> In turn, building ventilation rates matter for at least two major reasons: (a) they are related to public health and well being<sup>3,4</sup> and (b) they contribute substantially to energy use in buildings and consequently to total energy use.<sup>5,6</sup>

The most prominent gaseous effluent from humans is carbon dioxide (CO<sub>2</sub>) produced metabolically and emitted at rates of tens of grams per hour. The carbon dioxide level in an occupied indoor space is a proxy for the effectiveness of ventilation and has been found to associate with adverse health and well-being outcomes.<sup>7</sup> It had been long assumed that the cause of these adverse outcomes was not CO<sub>2</sub> itself, but rather some other as-yet-uncharacterized bioeffluent emissions whose indoor abundance would correlate with the metabolic CO<sub>2</sub> level. Recent studies have explored whether or not carbon dioxide is a direct-

acting indoor pollutant. Satish et al.<sup>8</sup> and Allen et al.<sup>9</sup> have shown that exposure to moderate CO<sub>2</sub> levels (1000–2500 ppm) in the absence of other bioeffluents can impair certain attributes of decision making. However, Zhang et al.<sup>10</sup> found that exposure to CO<sub>2</sub> alone at levels up to 3000 ppm did not degrade perceived air quality, induce acute health symptoms, or cause cognitive performance degradation. Zhang et al. did report, though, that, “exposures to bioeffluents with CO<sub>2</sub> at 3000 ppm reduced perceived air quality; increased the intensity of reported headache, fatigue, sleepiness, and difficulty in thinking clearly” and impaired certain indicators of cognitive functioning.

Within the indoor environment research community, VOCs are well recognized as a broad class of contaminants that pose important concerns for occupant health and well-being. Although occupants are recognized as a VOC source, relatively little work has been reported to characterize the chemical composition or rates of occupant emissions. This point is illustrated in the detailed report by Wolkoff<sup>11</sup> on the sources of indoor VOCs, which makes only brief mention of human occupants. A recent review by Weschler<sup>12</sup> does highlight the many ways that occupants influence indoor air chemistry, including through their emissions of VOCs.

Received: August 30, 2016

Revised: November 5, 2016

Accepted: November 7, 2016

Published: November 7, 2016

From other fields of study, there are substantial emerging literatures that aim to characterize VOCs associated with the human body and their components. A recent review tabulated 1840 VOCs associated with “breath, saliva, blood, milk, skin secretions, urine, and faeces in apparently healthy individuals.”<sup>13</sup> That review was motivated by interest in using chemical characterization of VOCs, for example, in exhaled breath, as a potential aid for the medical diagnosis of disease.<sup>14</sup> Another motivation for characterizing VOC emissions from humans is to assist with rescue operations in emergencies such as the aftermath of a building collapse.<sup>15–17</sup> These studies provide interesting information relevant to understanding occupant emissions of VOCs to indoor spaces. However, the focus for indoor environmental quality is not just to identify emitted species, but also to characterize rates of emissions. Furthermore, the emphasis on different components of the body is less pertinent than knowledge about the totality of emissions from building occupants, including those that originate from their metabolism, from personal care products that they have used, from their clothing, from chemical reactions occurring on their skin, and from the microbial communities that they host.

A few recent studies are directly relevant to the interest of characterizing occupant VOC emission rates. Veres et al.<sup>18</sup> reported on the air quality impacts of humans in a soccer stadium. Even for an open-roofed arena, the influence of the 31 000 attendees and their associated activities could be detected and quantified for several chemical analytes. Noteworthy are the emission rates (normalized to exhaled carbon dioxide) of ethanol, acetone, isoprene, 6-methyl-5-hepten-2-one (6-MHO), and decanal. Elevated ethanol emissions were associated with heavy beer consumption. The 6-MHO and decanal emissions were associated with ozone reactions with human skin oils. Clear signals of human occupancy could be seen and quantified in this environment even though the increment of metabolic carbon dioxide—about 80 ppm above ambient levels—indicates a much higher per-person effective ventilation rate than commonly applies to indoor spaces that are densely occupied. Williams et al.<sup>19</sup> found that human emission rates of certain volatile organic compounds varied according to audiovisual stimuli in the cinema. These interesting studies reveal information about VOC emissions in highly occupied spaces, but do not capture directly the circumstances that dominate for indoor occupancy, such as being in one’s own residence, being at work in an office, or being at school.<sup>20</sup>

Two studies have characterized human emissions of VOCs in university classrooms. In seminal work, Wang<sup>21</sup> applied time-integrated sorbent sampling with analysis by gas chromatography to characterize the emissions rate from university students of several compounds: acetone, acetaldehyde, allyl alcohol, acetic acid, amyl alcohol, butanoic acid, diethyl ketone, ethyl acetate, ethyl alcohol, methanol, phenol, and toluene. Analytical methods available for characterizing emissions have improved markedly in the four decades since Wang undertook his research. Liu et al.<sup>22</sup> monitored in real time the VOC levels in university classroom using a proton-transfer-reaction mass spectrometer. They utilized positive matrix factorization to detect a “human influence” component that varied with level of occupancy and with ventilation in a manner analogous to CO<sub>2</sub>. They reported that this component made an “average contribution of 40% to the measured daytime VOC concentration.”

To add new knowledge about the emission rates of VOCs from human occupants in an ordinary indoor environment, we conducted an intensive sampling campaign in a university classroom. Carried out over a two-week period, we made near continuous, time-resolved and chemically differentiated measurements of the broadest suite of VOCs that has been technologically possible thus far. We sampled from both the air supply and the classroom air and we monitored during both occupied and unoccupied periods. The application of a material balance model allows us to extract from the measured concentrations the chemical-specific net effective source rate entering the classroom air. The combination of sampling times and locations permits an apportionment of these source rates into contributions from occupants, indoor sources not related to occupancy, and outdoor air supplied by the ventilation system. For those chemicals substantially associated with occupancy, we are able to determine a per-occupant emission factor by combining the occupancy-associated aggregate emission rates with direct observations of the time-varying level of classroom occupancy.


The preliminary assessment of data from this field campaign revealed an unexpectedly strong contribution from cyclic volatile methylsiloxanes (cVMS). We reported briefly on those specific findings.<sup>23</sup> In this paper, we report on the other major results from this campaign, focusing on the emission rates of VOCs from human occupants of a university classroom.

## ■ MATERIALS AND METHODS

**Experimental Approach.** Air sampling was carried out in a normally functioning classroom (volume = 670 m<sup>3</sup>; air-exchange rate = 5 ± 0.5 h<sup>-1</sup>; mechanically ventilated, without recirculation, during the hours 8:00–20:45) at the University of California, Berkeley, California. Previous characterization of the classroom suggested negligible infiltration of air from other parts of the building because of the single-pass ventilation system and the absence of windows or exterior doors.<sup>24</sup> Details of the sampling strategy have been described.<sup>23</sup> Briefly, the monitoring phase was conducted over a two-week period in November 2014, including periods of stable occupancy for 19 separate class periods on five weekdays during which at least 17 occupants were in the classroom. One class period was excluded due to limited data availability, so results in this paper include 18 of the 19 class periods.

We continuously monitored concentrations of chemically differentiated VOCs, along with CO<sub>2</sub> and O<sub>3</sub> in the classroom air and in the ventilation air supplied to the classroom (supply air). A Teflon solenoid three-way valve enabled the sampling of the classroom air and supply air alternating at 5 min intervals. From each interval, the first 2 min of data were excluded and the remaining three were averaged. The resulting processed data set includes time-series measurements with six points per hour for the supply air and the classroom air, with each point representing the average condition over three contiguous minutes.

**VOC Measurements.** Mixing ratios of VOCs (in parts per billion) were measured using a proton-transfer-reaction time-of-flight mass spectrometer (PTR-TOF-MS; PTR-TOF 8000, IONICON Analytik GmbH). The PTR-TOF-MS recorded the mass spectrum for mass-to-charge ratios (*m/z*) 30–500 at a rate of 1 Hz, using H<sub>3</sub>O<sup>+</sup> as the primary reagent ion. VOCs with proton affinities greater than that of water undergo proton transfer reactions with H<sub>3</sub>O<sup>+</sup> in the drift tube, and are detected



**Figure 1.** Time series of mixing ratios in the classroom (solid line with markers) and supply air (dashed line) for  $\text{CO}_2$  and some representative VOCs. Vertical dashed lines define the duration of each class period on 13 November 2014; the average number of occupants in each class is noted above the arrows in the top panel.

by the mass spectrometer. PTR-TOF-MS is highly sensitive to alkenes, aromatics, alcohols, aldehydes, ketones, acids, esters, ethers, and many other compounds, but alkanes generally have proton affinities lower than water and are therefore not detected efficiently using  $\text{H}_3\text{O}^+$ .<sup>25,26</sup> However, even in the  $\text{H}_3\text{O}^+$  mode, there is up to 5% presence of impurity ions (i.e.,  $\text{O}_2^+$ , and  $\text{NO}^+$ ), which are sensitive to alkanes;<sup>27,28</sup> therefore, it is possible that abundant episodes or bursts of alkane emissions may have contributed to the signals seen by our instrument. VOCs reported here include all those detectable by PTR-TOF-MS as deployed in the classroom.

The instrument was calibrated daily with two multi-component VOC gas standard mixtures, including a total of 22 compounds with their protonated parent ions corresponding to these  $m/z$  values (5 of which are in both mixtures): 45.033, 33.034, 42.034, 59.048, 63.027, 69.069, 71.049, 73.028, 79.054, 83.086, 87.081, 93.07, 99.081, 107.086, 121.101, 137.133, and 146.977. Each chemical in the standard gas was present at a level of 1 ppm and was dynamically diluted to three concentrations (3, 6, and 9 ppb) using zero air of similar humidity to that of classroom air.

#### Data Analysis: Material Balance, Source Rates, and Emission Factors.

The total mass supply rate of each species entering the classroom was computed by material balance, assuming that the species is conserved and that the room air is well mixed. This mass supply rate was computed for the stable occupancy duration of each class period by applying an integral material balance in which the total supply rate is balanced by the change in the room air abundance and the total removal by means of ventilation. Each VOC in the classroom air could have one or more sources. We interpreted the data with the goal of apportioning the source rate for each VOC into three categories: (a) supply air (primarily from outdoor air), (b) human occupants (including their belongings), and (c) indoor sources that are not occupancy related, that is, from the building materials and room furnishings. The contributions from category (c) were determined utilizing classroom measurements during unoccupied periods at the end of each day. For this purpose, we analyzed data from periods of duration 40 min to 1 h beginning at least 1 h after the room became vacant. This protocol ensured that these periods had minimal contributions from human occupants and consequently the differences between the classroom air and the

supply air are likely to be associated with emissions from indoor sources other than occupants. We assumed that this non-occupancy emission rate was constant so that the value determined from end-of-day sampling could also be applied during periods of occupancy. In summary, the apportionment utilized direct measurements of supply air concentrations and the room ventilation rate to assess the contributions from ventilation supply. Measurements of the differences between supply air and room air when the room was vacant were interpreted to determine the nonoccupant contribution of indoor sources. Measurements of the differences between supply air and room air when the room was occupied, after correction for nonoccupancy contributions, were assigned to occupancy-associated emissions.

Detailed calculation procedures are reported in the [Supporting Information](#). An emission factor (EF,  $\mu\text{g person}^{-1} \text{h}^{-1}$ ) for each human-emitted VOC was calculated by dividing the occupancy-associated source rate for a given class session by the average number of the occupants in the classroom during that class session.<sup>23</sup>

## RESULTS AND DISCUSSION

Considering the whole sampling campaign, more than 400 ions were detected in the classroom air and supply air by the PTR-TOF-MS, which were filtered for internal ions and reduced using an abundance threshold. Consequently, 220 ions had mixing ratios (averaged over occupied periods for each sampling day) above 10 ppt in the classroom air and were evaluated further. Almost all of these 220 ions were on average more abundant in the classroom air than in the supply air. We focus on these 220 ions in this paper.

The detected chemicals were categorized into hydrocarbons ( $\text{C}_x\text{H}_y$ ); oxygenated organic compounds with 1 or 2 oxygen atoms in the molecule ( $\text{C}_x\text{H}_y\text{O}$ ,  $\text{C}_x\text{H}_y\text{O}_2$ ) such as carbonyls, alcohols, ethers, acids, diols, dicarbonyls, hydroxyl carbonyls and esters; and nitrogen (N)-, sulfur (S)- and silicon (Si)-containing organic compounds.


**Temporal Variation of Indoor VOCs.** Indoor VOC mixing ratios are expected to change temporally with ventilation, emission or uptake by indoor materials and humans. The temporal patterns of some representative VOCs are discussed here to provide evidence for their contributions to the composition of classroom air. The time series of  $\text{CO}_2$  mixing ratio in the classroom and supply air (top panel of [Figure 1](#)) serves as an independent tracer providing evidence for changes associated with ventilation and occupancy. Before the ventilation system was turned on at 8 AM, air that remained overnight in the classroom had the same stable  $\text{CO}_2$  mixing ratio as air in the supply duct. Following the beginning of ventilation system operation and occupancy (which occurred almost simultaneously), the  $\text{CO}_2$  level in the classroom is clearly higher than in the supply air and varies with the average number of occupants in the room during each class session (as labeled above the arrows in [Figure 1](#)). The average production rate of  $\text{CO}_2$  from occupants during all class periods was stable at  $21 \pm 3 \text{ g h}^{-1} \text{ person}^{-1}$ .<sup>20</sup> Consequently, and because the ventilation rate is constant when the mechanical ventilation system is operating, the temporal pattern of indoor  $\text{CO}_2$  level above the comparatively steady contribution of  $\text{CO}_2$  from supply air corresponds well with the number of occupants in each class session.

For VOCs primarily emitted from human metabolism, one expects a pattern similar to elevated classroom  $\text{CO}_2$ . The time

series for  $\text{C}_3\text{H}_6\text{O}$  (acetone) and  $\text{C}_3\text{H}_8$  (isoprene) mixing ratios (ppb) clearly show this pattern (second panel of [Figure 1](#)). When the classroom was occupied, human occupants contributed the dominant proportion of the mixing ratios in indoor air for these analytes. In a review on volatile emissions from healthy humans, isoprene and acetone were identified as the two most abundant organic components from human breath,<sup>13</sup> and so it is expected that they would be enriched in a densely occupied indoor environment. The elevated concentrations of acetone in the early morning before the ventilation was turned on were likely caused by janitorial staff cleaning the room; similar trends were observed for other chemicals that are expected to be present in cleaning products, such as monoterpenes.

The reaction of ozone with squalene found in skin oil produces 6-methyl-5-hepten-2-one (6-MHO) and 4-oxopentanal (4-OPA) as first- and second-generation products, respectively.<sup>29,30</sup> These compounds have been reported in connection to ozone-initiated chemistry on the human envelope in simulated aircraft cabin,<sup>31,32</sup> in offices,<sup>30,33</sup> and in classrooms.<sup>34</sup> The third frame of [Figure 1](#) displays the time-series of 6-MHO and 4-OPA concentrations measured in this study, indicating elevated levels associated with classroom occupancy (6-MHO 0.2–0.6 ppb, and 4-OPA 0.2–0.4 ppb) consistent with the reported values in Fischer et al.<sup>34</sup> of 0.2–0.7 ppb, and 0.12 ppb, respectively. The elevated 4-OPA in the morning, before the start of ventilation, also behaved similarly to the last measurement in Fischer et al.,<sup>34</sup> when that classroom remained empty for more than 1 h at the end of the day. Indoor sources in the absence of human occupants are likely to occur from continued ozone reactions with skin oils remaining on furniture and with shed skin flakes. Desorption from surfaces where the compounds had accumulated may also have contributed. In addition to 6-MHO and 4-OPA, several other gas-phase products of ozonolysis of human skin lipids, as reported in previous studies, were detected in the classroom air at significantly lower concentrations, including geranyl acetone ( $\text{C}_{13}\text{H}_{22}\text{OH}^+$ ), hydroxyacetone ( $\text{C}_3\text{H}_6\text{O}_2\text{H}^+$ , which may also be propionic acid), and 1,4-butanediol ( $\text{C}_4\text{H}_8\text{O}_2\text{H}^+$ ), along with minor products like 5-hydroxy-4-oxopentanal ( $\text{C}_5\text{H}_8\text{O}_3\text{H}^+$ ) and/or its isomer.<sup>29,31,35,36</sup> Ozone loss was observed in the classroom concurrent with increases of the ozone reaction products. Of these chemicals, 6-MHO, 4-OPA, and geranyl acetone exhibited contributions from indoor sources that were much larger than from the supply air. Hydroxyacetone and 1,4-butanediol, on the other hand, had similar levels of contribution from the supply air and from indoor sources.

A few VOCs such as monoterpenes ( $\text{C}_{10}\text{H}_{16}$  detected at  $m/z$  137.132) were observed to have large episodic increases associated with occupant activity, elevated above their already consistently higher mixing ratios in the classroom than in supply air (0.1–0.3 ppb). Monoterpenes are well-known biogenic VOCs, emitted by plants and fruits<sup>37,38</sup> and used in fragrances in personal care and cleaning products.<sup>39</sup> As illustrated in the fourth panel of [Figure 1](#), a short-term increase was observed in monoterpene level from about 2 to 20 ppb that persisted for about 10 min during the class of 9:40–11:00, followed by steady decay to the normal occupied-condition concentration. The strong episodic increase corresponds to a release of  $\sim 70$  mg of monoterpene and must have been caused by an occupant activity, for example, peeling an orange or applying a scented personal-care product. Based on high correlation ( $r > 0.96$ ) with  $m/z$  153.13 (citral) and  $m/z$  139.14



**Figure 2.** Median (shaded bars) and mean (circles) values of the VOC source rate ( $\text{mg h}^{-1}$ ) measured in 18 class sessions. The mass-to-charge ratio ( $m/z$ ) and assigned chemical formulas or names for the 20 most abundant ion/VOC groups are listed on the left of the graph. Of the remaining ions, those with known formulas are grouped by chemical composition and plotted at the bottom of the graph. The contribution of these top 20 individual ions and ion groups to the total source rate (mean value of 18 class periods) is shown in the pie chart on the right; ions with no empirical formula are summed and reported as “other”.


(methylisopropylcyclohexene), the ions typically found from citrus peel, and the lack of correlation with  $m/z$  155.14 (linalool), frequently present in perfumed products, the fruit-associated source seems more likely. We do not have records of occupant activities to confirm the specific source, but it clearly happened when students entered the classroom at the beginning of a lecture period. In contrast, the elevated levels of monoterpenes before ventilation was turned on in the morning were smaller, and presumably associated with cleaning activities or outgassing from indoor sources overnight.

The ventilation system supplies outdoor air and its associated VOCs through the building ducts to the classroom. Benzene ( $m/z$  79.053) and C8 aromatics ( $\text{C}_8\text{H}_{10}\text{H}^+$ ,  $m/z$  107.085) are common VOCs associated with gasoline vapors and with the exhaust of internal combustion engines. The classroom concentration of benzene and C8 aromatics clearly followed the pattern of supply air concentration, with some additional indoor source (not correlated with human occupancy) increasing the classroom-air mixing ratios in the case of benzene (lowest frame of Figure 1). Especially for the C8 aromatics, the time series of supply air and room air concentrations are nearly coincident after accounting for the

characteristic 12 min response time of the classroom air concentration to a change in supply air levels. (Recall that the classroom air-exchange rate is  $5 \text{ h}^{-1}$ ; the characteristic response time of indoor pollutant levels to a sudden change in outdoor levels is the reciprocal,  $0.2 \text{ h} = 12 \text{ min}$ .) The classroom concentration of benzene was constantly higher than the supply air level, providing evidence of a continuously emitting indoor nonoccupant source (e.g., from building materials or furnishings) in addition to the supply air source.

**VOC Source Rate and Its Apportionment.** To quantitatively evaluate the emission sources for indoor VOCs observable by PTR-TOF-MS, we calculated the source rates of the 220 VOCs/ions for each class period during the time of stable occupancy. The median total observed source rate of VOCs to the classroom during the 18 occupied periods was  $580 \text{ mg h}^{-1}$ . Occupant related emissions were the dominant source (57%), followed by supply air (35%), and indoor nonoccupant emissions (8%).

Quantitative apportionments of the three emission source categories are indicated in Figure 2 for the 20 most abundant VOCs/ions (referred to as VOCs hereafter). Source rates for the remaining ions for which chemical formulas could be



**Figure 3.** Human occupant emission factors (EF,  $\mu\text{g p}^{-1} \text{h}^{-1}$ ) for chemical species shown as relative contributions to the total (pie chart). The 20 most emitted compounds are specifically labeled, 62 (28% of total 220 ions) ions with no assigned formulas are grouped as “other” and the remaining ions are categorized by number of oxygen atoms and presence of nitrogen or sulfur in the molecule as  $\text{C}_x\text{H}_y$ ,  $\text{C}_x\text{H}_y\text{O}$ ,  $\text{C}_x\text{H}_y\text{O}_2$ ,  $\text{C}_x\text{H}_y\text{O}_3$ , N-containing and S-containing. The values of EF for the listed compounds and ion groups are shown on the right.

confidently assigned are summarized according to chemical composition families ( $\text{C}_x\text{H}_y$ ,  $\text{C}_x\text{H}_y\text{O}$ ,  $\text{C}_x\text{H}_y\text{O}_2$ , etc.). Observed ions for which empirical chemical formulas could not be confidently assigned were summed and reported as “other”; these accounted for just 5% of the total mass, as shown in the pie chart. Detailed results for all observed ions are summarized in Table S1 in the [Supporting Information](#).

The six most abundant VOCs accounted for 62% of the total measured source rate (shown in the top group of [Figure 2](#)), and the 20 most abundant VOCs accounted for 80% of the total. The label cVMS represents the sum of four cyclic siloxane compounds, including D3 (hexamethylcyclotrisiloxane), D4 (octamethylcyclotetrasiloxane), D5 (decamethylcyclopentasiloxane) and D6 (dodecamethylcyclohexasiloxane), as we have previously reported.<sup>23</sup> The dominant cVMS observed was D5.

Comparison of the median (shaded bars) and mean (circles) source rates is indicative of the level of variance in sources during the stably occupied periods. A mean emission value higher than the median indicates variability in source rates among class sessions along with positive skewness. About half of the VOCs among the top 20 were observed to have highly variable sources, but the sources may be variable for different reasons. Among the contributors to variability are time-varying emissions from occupants (e.g., the D5 emission factor declined with time of day), varying levels of urban air pollution (C8 aromatics and toluene), or variability in the occurrence of specific short-term emissions events (e.g., peeling citrus fruit that releases monoterpenes).

While indoor nonoccupant emission sources made relatively small contributions to the total (8%), they do provide discernible contributions to some of the top 20 VOCs, mainly

among organic acids (acetic acid, formic acid, acid fragment  $m/z$  43.018), ketones (acetone), alcohols (methanol, ethanol), aldehydes (acetaldehyde, hexanal), and other oxygenated VOCs ( $\text{C}_9\text{H}_{10}\text{OH}^+$ ). Other than the observation that indoor emissions of formaldehyde were low in this classroom, these findings are consistent with the literature regarding indoor sources from building materials.<sup>40,41</sup> Emissions of these chemicals from nonoccupant indoor sources have been extensively characterized in past studies, so we do not probe more deeply here.

The contributions from supply air were a typical mixture of outdoor air VOCs expected to be found in urban areas such as acetone, acetic acid, methanol, acetaldehyde, monoterpenes, organic acids, isoprene, ethanol, and components of gasoline, such as benzene, toluene, and C8 aromatics, etc.<sup>42,43</sup> In this study, the mean contributions of these outdoor air VOCs to classroom air was 35% of the total. That level of contribution could vary substantially in other areas, depending on the level of outdoor air pollution and the building ventilation rates. A more detailed analysis of the composition and source characteristics of outdoor air in this study would not provide general results relevant to predicting indoor air concentrations elsewhere.

A noteworthy result from this study is that human occupant emissions were the dominant source of VOC during classroom-occupied periods, contributing 57% of the total mass of quantified sources. Of the top 20 VOC, all but three (C8 aromatics, toluene, and  $\text{C}_4\text{H}_6\text{O}$ ) had substantial contributions from human occupants. Three cVMS (primarily D5, along with D4 and D6) together comprised  $\sim 1/3$  of the total indoor VOC mass concentration in the classroom and were predominantly associated with occupant emissions.<sup>23</sup> Other prominent VOCs

whose source was  $\sim 1/3$  or more from human occupants included acetone, isoprene, acetic acid, methanol, acetaldehyde, monoterpenes, organic acids (formic, acid fragments), ethanol, hexanal, (iso) butyl and (iso) propyl fragments, hydroxyacetone, and the products of ozone reactions with skin oil, that is, 4-OPA and 6-MHO. Hundreds of organic acids have been previously detected in volatiles from human skin secretions,<sup>13,44,45</sup> including acetic and formic acid. Previous studies using PTR-TOF-MS to analyze human breath show most of the same dominant 15 compounds including acetone, methanol, acetaldehyde, ethanol, formic acid, (iso) butyl fragment ( $C_4H_8H^+$ ), and isoprene.<sup>46</sup> One significant difference worth noting is that Herbig et al. reported  $m/z$  71.049 ( $C_4H_6OH^+$ ) in breath, but we observed no occupancy-related source and instead observed indoor nonoccupant sources of this ion to be dominant. The  $C_4H_6OH^+$  ion could be from methyl vinyl ketone or from methacrolein,<sup>47</sup> which are products of the atmospheric oxidation of isoprene.

Observed occupant emissions included N- and S-containing VOCs. Volatile sulfur compounds, which are the main cause for oral malodor,<sup>48–50</sup> and which contribute to underarm odors,<sup>44</sup> were detected in classroom air. Examples include methanethiol ( $m/z$  49.013,  $CH_3SH^+$ ) and dimethyl sulfide ( $m/z$  63.025,  $C_2H_6SH^+$ ). A large number of nitrogen-containing compounds have also been identified as volatiles from human body, mostly in human breath,<sup>13</sup> consistent with our observations in classroom air.

**Emission Factors for Human-Emitted Compounds.** As might be expected, we found that the occupant-averaged  $CO_2$  generation rate was relatively stable across all class periods, indicating similar average metabolic activity levels among occupants. Analogous characterization of the spectrum of occupant-emitted VOCs was conducted by calculating the per-person emission factors (EFs,  $\mu g\ p^{-1}\ h^{-1}$ ), as listed in Figure 3, with a pie chart indicating percent mass contributions for each species to the total human occupant emission rates.

The cVMS had the highest EF among all occupancy-associated VOC emissions.<sup>23</sup> The next highest EF values were for acetone, acetic acid, monoterpenes, isoprene, methanol, acetaldehyde, ethanol, formic, and other acid fragments. Wang<sup>21</sup> reported subject-weighted emission factors in a university classroom for some of the same organic compounds that we observed. The EFs reported by Wang for chemicals known to be major metabolic products in human breath, including acetone, acetaldehyde, acetic acid, methanol, and ethanol, were all within a factor of 2 of our measured EFs. This agreement is good for studies using completely independent populations, measurement techniques, standards, and undertaken four decades apart. However, surprisingly, the EFs Wang reported for toluene and phenol were 1–2 orders of magnitude higher than the values we obtained (308 and 396  $\mu g\ p^{-1}\ h^{-1}$  versus 6 and 12  $\mu g\ p^{-1}\ h^{-1}$ , respectively). Human exposure to toluene in many urban areas has been dramatically reduced over the past 40 years as a result of improved emission controls for outdoor air pollutants. We might also speculate about another potential factor contributing to the differences observed: a reduction over time in the use of aromatic compounds, which have been gradually withdrawn from consumer products and from other items that would be carried to class by students.

The total average VOC emission factor for human occupants that we have obtained, 6.3  $mg\ h^{-1}\ person^{-1}$ , is only partly accounted for by the VOCs reported in past studies. As shown

in Figure 3, cVMS accounted for 44%, acetone for 16%, the sum of identified acids, acid fragments, methanol, ethanol, isoprene, and monoterpenes accounted for another  $\sim 20\%$ , and the remaining  $\sim 20\%$  consisted of a large array of VOCs with small but measurable EFs summed into classes of chemicals by their elemental composition ( $C_xH_y$ ,  $C_xH_yO$ ,  $C_xH_yO_2$ ,  $C_xH_yO_3$ , N-containing, S-containing). (See Table S2 for a list of VOCs for which the occupancy-associated emissions were greater than 25% of the total source strength.) Only a few percent of the total observed EF are unidentified by chemical formula owing to a lack of any definitive match to the observed exact ion mass. While some of the most abundant chemicals had been previously reported, the full range of reported species and the time resolution of our measurements provide novel contributions to knowledge regarding human influence on the composition of indoor air. The specific population studied here (engineering students in a university classroom in northern California), cannot be assumed to be representative of the broader human population. Additional research focusing on emissions from different groups of people in different indoor spaces are likely to reveal important variability in occupant emission rates due to age, activity, health status, emotional state,<sup>19</sup> or other factors.

In this study, we have examined the full spectrum of VOCs emitted indoors in a university classroom and found that human occupants were the major contributor to the mass of indoor VOCs, exceeding contributions from supply air and from indoor nonoccupant sources. The per-person emission factors for human-emitted chemicals have rarely been reported in previous studies. Our findings improve knowledge of human emissions influencing the chemistry of indoor environments, including chemically speciated per-capita emission factors that can be used for modeling indoor air quality.

## ■ ASSOCIATED CONTENT

### 📄 Supporting Information

Sampling Site and Strategy; Integral Material Balance for Evaluating VOC Source Rates and its Apportionment; PTR-TOF-MS VOC Measurement Details Table S1. Source rates ( $mg\ h^{-1}$ ) for 220 ions measured in all stable class periods during 5 sampling days. Table S2. Emission factor ( $\mu g\ h^{-1}\ p^{-1}$ ) for VOCs with occupant-source contributing  $>25\%$  of the total median source rates. The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.est.6b04415.

(PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*Phone: (510) 643-2451; e-mail: [ahg@berkeley.edu](mailto:ahg@berkeley.edu).

### ORCID

Xiaochen Tang: 0000-0003-4168-9871

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

This work was funded by the Alfred P. Sloan Foundation via Grant 2013-10-04. We thank Seema Bhangar and Robin Weber for technical assistance.

## REFERENCES

- (1) Yaglou, C. P.; Riley, E. C.; Coggins, D. I. Ventilation requirements. *ASHVE Trans.* **1936**, *42*, 133–162.
- (2) Persily, A. Challenges in developing ventilation and indoor air quality standards: The story of ASHRAE Standard 62. *Build. Environ.* **2015**, *91*, 61–69.
- (3) Sundell, J.; Levin, H.; Nazaroff, W. W.; Cain, W. S.; Fisk, W. J.; Grimsrud, D. T.; Gyntelberg, F.; Li, Y.; Persily, A. K.; Pickering, A. C.; Samet, J. M.; Spengler, J. D.; Taylor, S. T.; Weschler, C. J. Ventilation rates and health: Multidisciplinary review of the scientific literature. *Indoor Air* **2011**, *21*, 191–204.
- (4) Carrer, P.; Wargocki, P.; Fanetti, A.; Bischof, W.; de Oliveira Fernandes, E.; Hartmann, T.; Kephelopoulou, S.; Palkonen, S.; Seppänen, O. What does the scientific literature tell us about the ventilation–health relationship in public and residential buildings? *Build. Environ.* **2015**, *94*, 273–286.
- (5) Fisk, W. J.; Black, D.; Brunner, G. Changing ventilation rates in U.S. offices: Implications for health, work performance, energy, and associated economics. *Build. Environ.* **2012**, *47*, 368–372.
- (6) Chenari, B.; Carrilho, J. D.; da Silva, M. G. Towards sustainable, energy-efficient and healthy ventilation strategies in buildings: A review. *Renewable Sustainable Energy Rev.* **2016**, *59*, 1426–1447.
- (7) Seppänen, O. A.; Fisk, W. J.; Mendell, M. J. Association of ventilation rates and CO<sub>2</sub> concentrations with health and other responses in commercial and institutional buildings. *Indoor Air* **1999**, *9*, 226–252.
- (8) Satish, U.; Mendell, M. J.; Shekhar, K.; Hotchi, T.; Sullivan, D.; Streufert, S.; Fisk, W. J. Is CO<sub>2</sub> an indoor pollutant? Direct effects of low-to-moderate CO<sub>2</sub> concentrations on human decision-making performance. *Environ. Health Perspect.* **2012**, *120*, 1671–1677.
- (9) Allen, J. G.; MacNaughton, P.; Satish, U.; Santanam, S.; Vallarino, J.; Spengler, J. D. A controlled exposure study of green and conventional office environments. *Environ. Health Perspect.* **2016**, *124* (6), 805–812.
- (10) Zhang, X.; Wargocki, P.; Lian, Z.; Thyregod, C. Effects of exposure to carbon dioxide and bioeffluents on perceived air quality, self-assessed acute health symptoms, and cognitive performance. *Indoor Air* **2016**, n/a.
- (11) Wolkoff, P. Volatile organic compounds – Sources, measurements, emissions, and the impact on indoor air quality. *Indoor Air* **1995**, *5* (Suppl. 3), 1–73.
- (12) Weschler, C. J. Roles of the human occupant in indoor chemistry. *Indoor Air* **2016**, *26*, 6–24.
- (13) de Lacy Costello, B.; Amann, A.; Al-Kateb, H.; Flynn, C.; Filipiak, W.; Khalid, T.; Osborne, D.; Ratcliffe, N. M. A review of the volatiles from the healthy human body. *J. Breath Res.* **2014**, *8*, 014001.
- (14) Amann, A.; Miekisch, W.; Schubert, J.; Buszewski, B.; Ligor, T.; Jezierski, T.; Pleil, J.; Risby, T. Analysis of exhaled breath for disease detection. *Annu. Rev. Anal. Chem.* **2014**, *7*, 455–482.
- (15) Mochalski, P.; Unterkofler, K.; Teschl, G.; Amann, A. Potential of volatile organic compounds as markers of entrapped humans for use in urban search-and-rescue operations. *TrAC, Trends Anal. Chem.* **2015**, *68*, 88–106.
- (16) Huo, R.; Agapiou, A.; Bocos-Bintintan, V.; Brown, L. J.; Burns, C.; Creaser, C. S.; Devenport, N. A.; Gao-Lau, B.; Guallar-Hoyas, C.; Hildebrand, L.; Malkar, A.; Martin, H. J.; Moll, V. H.; Patel, P.; Ratiu, A.; Reynolds, J. C.; Sielemann, S.; Slodzynski, R.; Statheropoulos, M.; Turner, M. A.; Vautz, W.; Wright, V. E.; Thomas, C. L. P. The trapped human experiment. *J. Breath Res.* **2011**, *5*, 046006.
- (17) Vautz, W.; Slodzynski, R.; Hariharan, C.; Seifert, L.; Nolte, J.; Fobbe, R.; Sielemann, S.; Lao, B. C.; Huo, R.; Thomas, C. L. P.; Hildebrand, L. Detection of metabolites of trapped humans using ion mobility spectrometry coupled with gas chromatography. *Anal. Chem.* **2013**, *85*, 2135–2142.
- (18) Veres, P. R.; Faber, P.; Drownick, F.; Lelieveld, J.; Williams, J. Anthropogenic sources of VOC in a football stadium: Assessing human emissions in the atmosphere. *Atmos. Environ.* **2013**, *77*, 1052–1059.
- (19) Williams, J. R.; Stöner, C.; Wicker, J.; Krauter, N.; Derstroff, B.; Bourtsoukidis, E.; Klüpfel, T.; Kramer, S. Cinema audiences reproducibly vary the chemical composition of air during films, by broadcasting scene specific emissions on breath. *Sci. Rep.* **2016**, *6*, 25464.
- (20) Klepeis, N. E.; Nelson, W. C.; Ott, W. R.; Robinson, J. P.; Tsang, A. M.; Switzer, P.; Behar, J. V.; Hern, S. C.; Engelmann, W. H. The National Human Activity Pattern Survey (NHAPS): A resource for assessing exposure to environmental pollutants. *J. Exposure Anal. Environ. Epidemiol.* **2001**, *11*, 231–252.
- (21) Wang, T. C. A study of bioeffluents in a college classroom. *ASHRAE Transactions* **1975**, *81* (I), 32–44.
- (22) Liu, S.; Li, R.; Wild, R. J.; Warneke, C.; de Gouw, J. A.; Brown, S. S.; Miller, S. L.; Luongo, J. C.; Jimenez, J. L.; Ziemann, P. J. Contribution of human-related sources to indoor volatile organic compounds in a university classroom. *Indoor Air* **2016**, *26*, 925–938.
- (23) Tang, X.; Misztal, P. K.; Nazaroff, W. W.; Goldstein, A. H. Siloxanes are the most abundant volatile organic compound emitted from engineering students in a classroom. *Environ. Sci. Technol. Lett.* **2015**, *2*, 303–307.
- (24) Bhangar, S.; Huffman, J. A.; Nazaroff, W. W. Size-resolved fluorescent biological aerosol particle concentrations and occupant emissions in a university classroom. *Indoor Air* **2014**, *24*, 604–617.
- (25) Gueneron, M.; Erickson, M. H.; VanderSchelden, G. S.; Jobson, B. T. PTR-MS fragmentation patterns of gasoline hydrocarbons. *Int. J. Mass Spectrom.* **2015**, *379*, 97–109.
- (26) Erickson, M. H.; Gueneron, M.; Jobson, T. Measuring long chain alkanes in diesel engine exhaust by thermal desorption PTR-MS. *Atmos. Meas. Tech.* **2014**, *7*, 225–239.
- (27) Amador Muñoz, O.; Misztal, P. K.; Weber, R.; Worton, D. R.; Zhang, H.; Drozd, G.; Goldstein, A. H. Sensitive detection of *n*-alkanes using a mixed ionization mode proton-transfer-reaction mass spectrometer. *Atmos. Meas. Tech.* **2016**, *9*, 5315–5329.
- (28) Koss, A. R.; Warneke, C.; Yuan, B.; Coggon, M. M.; Veres, P. R.; de Gouw, J. A. Evaluation of NO<sup>+</sup> reagent ion chemistry for online measurements of atmospheric volatile organic compounds. *Atmos. Meas. Tech.* **2016**, *9*, 2909–2925.
- (29) Fruekilde, P.; Hjorth, J.; Jensen, N. R.; Kotzias, D.; Larsen, B. Ozonolysis at vegetation surfaces: A source of acetone, 4-oxopentanal, 6-methyl-5-hepten-2-one, and geranyl acetone in the troposphere. *Atmos. Environ.* **1998**, *32*, 1893–1902.
- (30) Wisthaler, A.; Weschler, C. J. Reactions of ozone with human skin lipids: Sources of carbonyls, dicarbonyls, and hydroxycarbonyls in indoor air. *Proc. Natl. Acad. Sci. U. S. A.* **2010**, *107*, 6568–6575.
- (31) Wisthaler, A.; Tamás, G.; Wyon, D. P.; Strøm-Tejse, P.; Space, D.; Beauchamp, J.; Hansel, A.; Märk, T. D.; Weschler, C. J. Products of ozone-initiated chemistry in a simulated aircraft environment. *Environ. Sci. Technol.* **2005**, *39*, 4823–4832.
- (32) Weschler, C. J.; Wisthaler, A.; Cowlin, S.; Tamás, G.; Strøm-Tejse, P.; Hodgson, A. T.; Destailats, H.; Herrington, J.; Zhang, J. J.; Nazaroff, W. W. Ozone-initiated chemistry in an occupied simulated aircraft cabin. *Environ. Sci. Technol.* **2007**, *41*, 6177–6184.
- (33) Bakó-Biró, Z. Human Perception, SBS Symptoms and Performance of Office Work during Exposure to Air Polluted by Building Materials and Personal Computers. Ph. D. Thesis, Department of Mechanical Engineering, Technical University of Denmark, 2004.
- (34) Fischer, A.; Ljungström, E.; Langer, S. Ozone removal by occupants in a classroom. *Atmos. Environ.* **2013**, *81*, 11–17.
- (35) Zhang, Z. M.; Cai, J. J.; Ruan, G. H.; Li, G. K. The study of fingerprint characteristics of the emanations from human arm skin using the original sampling system by SPME-GC/MS. *J. Chromatogr. B* **2005**, *822*, 244–252.
- (36) Weisel, C.; Weschler, C. J.; Mohan, K.; Vallarino, J.; Spengler, J. D. Ozone and ozone byproducts in the cabins of commercial aircraft. *Environ. Sci. Technol.* **2013**, *47*, 4711–4717.
- (37) Vartiainen, E.; Kulmala, M.; Ruuskanen, T. M.; Taipale, R.; Rinne, J.; Vehkamäki, H. Formation and growth of indoor air aerosol



particles as a result of d-limonene oxidation. *Atmos. Environ.* **2006**, *40*, 7882–7892.

(38) Gentner, D. R.; Ormeño, E.; Fares, S.; Ford, T. B.; Weber, R.; Park, J.-H.; Brioude, J.; Angevine, W. M.; Karlik, J. F.; Goldstein, A. H. Emissions of terpenoids, benzenoids, and other biogenic gas-phase organic compounds from agricultural crops and their potential implications for air quality. *Atmos. Chem. Phys.* **2014**, *14*, 5393–5413.

(39) Corsi, R. L.; Siegel, J.; Karamalegos, A.; Simon, H.; Morrison, G. C. Personal reactive clouds: Introducing the concept of near-head chemistry. *Atmos. Environ.* **2007**, *41*, 3161–3165.

(40) Brown, S. K. Volatile organic pollutants in new and established buildings in Melbourne, Australia. *Indoor Air* **2002**, *12*, 55–63.

(41) Liang, W.; Yang, C.; Yang, X. Long-term concentrations of volatile organic compounds in a new apartment in Beijing, China. *Build. Environ.* **2014**, *82*, 693–701.

(42) Liu, W.; Zhang, J.; Zhang, L.; Turpin, B. J.; Weisel, C. P.; Morandi, M. T.; Stock, T. H.; Colome, S.; Korn, L. R. Estimating contributions of indoor and outdoor sources to indoor carbonyl concentrations in three urban areas of the United States. *Atmos. Environ.* **2006**, *40*, 2202–2214.

(43) Cheng, M.; Galbally, I. E.; Molloy, S. B.; Selleck, P. W.; Keywood, M. D.; Lawson, S. J.; Powell, J. C.; Gillett, R. W.; Dunne, E. Factors controlling volatile organic compounds in dwellings in Melbourne, Australia. *Indoor Air* **2016**, *26*, 219–230.

(44) Gallagher, M.; Wysocki, C. J.; Leyden, J. J.; Spielman, A. I.; Sun, X.; Preti, G. Analyses of volatile organic compounds from human skin. *Br. J. Dermatol.* **2008**, *159*, 780–791.

(45) Kuhn, F.; Natsch, A. Body odour of monozygotic human twins: A common pattern of odorant carboxylic acids released by bacterial aminoacylase from axilla secretions contributing to an inherited body odour type. *J. R. Soc. Interface* **2009**, *6*, 377–392.

(46) Herbig, J.; Müller, M.; Schallhart, S.; Titzmann, T.; Graus, M.; Hansel, A. On-line breath analysis with PTR-TOF. *J. Breath Res.* **2009**, *3*, 027004.

(47) de Gouw, J.; Warneke, C. Measurements of volatile organic compounds in the earth's atmosphere using proton-transfer-reaction mass spectrometry. *Mass Spectrom. Rev.* **2007**, *26*, 223–257.

(48) Tonzetich, J. Production and origin of oral malodor: a review of mechanisms and methods of analysis. *J. Periodontol.* **1977**, *48*, 13–20.

(49) Persson, S.; Edlund, M. B.; Claesson, R.; Carlsson, J. The formation of hydrogen sulfide and methyl mercaptan by oral bacteria. *Oral Microbiol. Immunol.* **1990**, *5*, 195–201.

(50) Miyazaki, H.; et al. Oral malodor in the general population of Japan. In *Bad Breath: Research Perspectives*; Rosenberg, M., Ed.; Ramot Publishing, Tel Aviv, Israel, 1995; pp 119–137.